

## Note

 $M_2$ -equipackable graphs<sup>☆</sup>

 Yuqin Zhang<sup>a</sup>, Yonghui Fan<sup>b</sup>
<sup>a</sup>Department of Mathematics, Beijing Institute of Technology, 100081, Beijing, China

<sup>b</sup>College of Mathematics, Hebei Normal University, 050016, Shijiazhuang, China

Received 25 November 2005; received in revised form 12 March 2006; accepted 27 March 2006

Available online 2 May 2006

## Abstract

Let  $H$  be a fixed graph. An  $H$ -packing of  $G$  is a set of edge disjoint subgraphs of  $G$  each isomorphic to  $H$ . An  $H$ -packing in  $G$  with  $k$  copies  $H_1, H_2, \dots, H_k$  of  $H$  is called *maximal* if  $G - \bigcup_{i=1}^k E(H_i)$  contains no subgraph isomorphic to  $H$ . An  $H$ -packing in  $G$  with  $k$  copies  $H_1, H_2, \dots, H_k$  of  $H$  is called *maximum* if no more than  $k$  edge disjoint copies of  $H$  can be packed into  $G$ . A graph  $G$  is called  *$H$ -equipackable* if every maximal  $H$ -packing in  $G$  is also a maximum  $H$ -packing in  $G$ . By  $M_t$ ,  $t \geq 1$ , we denote a matching having  $t$  edges. In this paper, we investigate the characterization of  $M_2$ -equipackable graphs.  
© 2006 Elsevier B.V. All rights reserved.

**Keywords:** Packing; Packable; Randomly packable; Equipackable; Matching

## 1. Preliminaries and known results

A vertex of degree 0 is called an isolated vertex (or an isolate). The graphs characterized in this paper do not have any isolated vertices. A graph  $G$  has order  $|V(G)|$  and size  $|E(G)|$ . The path and circuit on  $k$  vertices are denoted by  $P_k$  and  $C_k$ , respectively. A matching in the graph  $G$  is a set of independent edges in  $G$ . By  $M_t$ ,  $t \geq 1$ , we denote a matching having  $t$  edges. Let  $H$  be a subgraph of  $G$ . By  $G - H$ , we denote the graph remaining after we delete from  $G$  the edges of  $H$  and any resulting isolated vertices. A collection of edge disjoint copies of  $H$ , say  $H_1, H_2, \dots, H_k$ , where each  $H_i$  is a subgraph of  $G$ , is called an  $H$ -packing in  $G$ . An  $H$ -packing in  $G$  with  $k$  copies  $H_1, H_2, \dots, H_k$  of  $H$  is called *maximal* if  $G - \bigcup_{i=1}^k E(H_i)$  contains no subgraph isomorphic to  $H$ . An  $H$ -packing in  $G$  with  $k$  copies  $H_1, H_2, \dots, H_k$  of  $H$  is called *maximum* if no more than  $k$  edge disjoint copies of  $H$  can be packed into  $G$ . A graph  $G$  is called  *$H$ -packable* if there exists an  $H$ -packing of  $G$  which uses all edges in  $G$  and  $G$  is called *randomly  $H$ -packable* if every maximal  $H$ -packing in  $G$  uses all edges in  $G$ . There have been many results on  $H$ -packable graphs and randomly  $H$ -packable graphs (see [1,3]). The following results when  $H = M_2$  are very useful to our work:

**Theorem 1.1** (Caro [4]). *Let  $G$  be a graph of size  $2m > 0$  and without isolated vertices. Then  $G$  is  $M_2$ -packable if and only if  $\Delta(G) \leq m$  and  $G$  is not isomorphic to  $K_3 \cup K_2$ .*

<sup>☆</sup> This research was supported by National Natural Science Foundation of China (10571042), NSF of Hebei (A2005000144), Tianyuan Fund for Mathematics (10426013) and Fund of Hebei Province for Doctorate (B2004114).

E-mail addresses: [yqinzhang@163.com](mailto:yqinzhang@163.com), [yqinzhang@126.com](mailto:yqinzhang@126.com) (Y. Zhang).

**Theorem 1.2** (Ruiz [4]). Let  $G$  be a graph of size  $2m > 0$  without isolated vertices and suppose  $G$  is not isomorphic to  $M_2$ , then  $G$  is randomly  $M_2$ -packable if and only if  $G \in \mathcal{F}$ , where  $\mathcal{F} = \{K_4, C_4, 2K_3, K_3 \cup K_{1,3}\} \cup \{2mK_2, 2K_{1,m} | m \geq 2\}$ .

As a relaxation of random  $H$ -packability, Hartnell and Vestergaard [2] introduced the definition of  $H$ -equipackable. A graph  $G$  is called  $H$ -equipackable if every maximal  $H$ -packing in  $G$  is also a maximum  $H$ -packing in  $G$ . Hartnell and Vestergaard [2] first characterized the  $P_3$ -equipackable graphs of girth five or more. Later, Vestergaard [6] gave a characterization of  $P_3$ -equipackable graphs with all valences at least two. Recently, Vestergaard and a coauthor have extended the results of [2,6] by characterizing all  $P_3$ -equipackable graphs without any restrictions on the graph.

In this paper, we investigate  $M_2$ -equipackable graphs.

Note that when  $G$  is congruent to  $K_2$  or  $P_3$  there does not exist a copy of  $M_2$ . When  $G \cong M_2$ ,  $G$  is clearly  $M_2$ -equipackable. So, in the sections that follow, we consider graphs with size at least 3.

## 2. $M_2$ -equipackable graphs with even size

**Lemma 2.1.** Let  $G$  be a graph with size  $2m$  and maximum degree  $d$ . If  $d > m$ , then the number of  $M_2$  in the maximum  $M_2$ -packing of  $G$  is  $2m - d$ .

**Proof.** Assume that  $v$  is a vertex with degree  $d$  and  $v_1, v_2, \dots, v_d$  are the neighbors of  $v$ . Let  $E_1 = \{vv_i | i = 1, 2, \dots, d\}$  and  $E_2 = E(G) - E_1 = \{e_1, e_2, \dots, e_{2m-d}\}$ . Since  $d$  and  $m$  are both positive integers and  $d > m$ ,  $|E_1| - |E_2| = d - (2m - d) = 2(d - m) \geq 2$ . Clearly, each edge of  $E_2$  is adjacent to at most two edges of  $E_1$ , so for any  $e_i \in E_2$ , there exists an edge  $vv_j \in E_1$  such that  $\{e_i, vv_j\}$  forms a copy of  $M_2$ . Remove  $\{e_i, vv_j\}$ . Let  $E_1^{(1)} = E_1 - \{vv_j\}$  and  $E_2^{(1)} = E_2 - \{e_i\}$ . In the same way, each edge in  $E_2^{(1)}$  has at most two neighbors in  $E_1^{(1)}$ , then we can get another copy of  $M_2$  and remove it. Repeating this process  $2m - d$  times, we can remove all the edges of  $E_2$ , each of which along with one edge of  $E_1$  forms a copy of  $M_2$ . And the remaining edges in  $E_1$  contains no  $M_2$ . So the removed  $2m - d$  copies of  $M_2$  form a maximal  $M_2$ -packing of  $G$ . Because all edges in  $E_1$  are adjacent and no two edges in  $E_2$  belong to one copy of  $M_2$  at the same time, this  $M_2$ -packing is maximum.  $\square$

**Theorem 2.2.** Let  $G$  be a graph with size  $2m > 2$ , then  $G$  is  $M_2$ -equipackable if and only if  $G$  satisfies one of the following:

- (1)  $G \cong K_3 \cup K_2$ ;
- (2)  $G \in \mathcal{F}$ , where  $\mathcal{F} = \{K_4, C_4, 2K_3, K_3 \cup K_{1,3}\} \cup \{2mK_2, 2K_{1,m} | m \geq 2\}$ ;
- (3)  $\Delta(G) = d > m$ , and for any vertex  $v$  whose degree is  $d$ , the subgraph induced by  $E(G - v)$  must be  $K_{1,2m-d}$  or  $K_3$ .

**Proof.** We can easily verify that the graphs described in (1)–(3) are all  $M_2$ -equipackable.

Conversely, let  $G$  be an  $M_2$ -equipackable graph with size  $2m$ . We consider two cases:

Case 1:  $\Delta(G) \leq m$ .

Subcase 1: When  $G \cong K_3 \cup K_2$ ,  $G$  is clearly  $M_2$ -equipackable.

Subcase 2: When  $G$  is not isomorphic to  $K_3 \cup K_2$ , we have:

By Theorem 1.1,  $G$  is  $M_2$ -packable. So the number of  $M_2$  in the maximum  $M_2$ -packing is  $m$ . If  $G$  is not randomly  $M_2$ -packable, then there exists a maximal  $M_2$ -packing which does not use all edges in  $G$  and, consequently, which is not maximum. This contradicts the fact that  $G$  is  $M_2$ -equipackable. So  $G$  must be randomly  $M_2$ -packable. By Theorem 1.2,  $G \in \mathcal{F} = \{K_4, C_4, 2K_3, K_3 \cup K_{1,3}\} \cup \{2mK_2, 2K_{1,m} | m \geq 2\}$ .

Case 2:  $\Delta(G) = d > m$ .

By Lemma 2.1, the number of  $M_2$  in the maximum  $M_2$ -packing in  $G$  is  $2m - d$ . Let  $v$  be a vertex with maximum degree. If there are two edges (say  $\{e, f\}$ ) in  $G - v$  which are not adjacent, then after removing  $\{e, f\}$  (we denote  $G - \{e, f\}$  by  $G_1$ ),  $\Delta(G_1) = d > m > m - 1$ . The graph  $G_1$  also satisfies Lemma 2.1. So we can get a maximum  $M_2$ -packing in  $G_1$  with  $2(m - 1) - d$  copies of  $M_2$  which, along with  $\{e, f\}$ , form a maximal  $M_2$ -packing. Obviously this resulting maximal packing with only  $2m - d - 1$  copies of  $M_2$  is not maximum, which contradicts that  $G$  is

$M_2$ -equipackable. So all edges in  $G - v$  are mutually adjacent; that is, the subgraph induced by  $E(G - v)$  must be  $K_{1,2m-d}$  or  $K_3$ .  $\square$

### 3. $M_2$ -equipackable graphs with odd size

The following lemma can be proven in a manner similar to Lemma 2.1. The proof is thus omitted.

**Lemma 3.1.** *Let  $G$  be a graph with size  $2m + 1$ . If  $\Delta(G) = d > m + 1$ , then the number of  $M_2$  in the maximum  $M_2$ -packing of  $G$  is  $2m + 1 - d$ .*

**Theorem 3.2.** *Let  $G$  be a graph with size  $2m + 1$  and  $\Delta(G) = d \geq m + 2$ , then  $G$  is  $M_2$ -equipackable if and only if, for any vertex  $v$  whose degree is  $d$ , the subgraph induced by  $E(G - v)$  must be  $K_{1,2m+1-d}$  or  $K_3$ .*

**Proof.** Applying the same technique as in the proof of Case 2 of Theorem 2.2, we can easily get the result.  $\square$

**Lemma 3.3.** *Let  $G$  be a graph with size  $2m + 1$  and  $F$  be a maximal  $M_2$ -packing which omits all the edges of a star  $K_{1,3}$ . If  $F$  contains a copy (say  $\{e, f\}$ ) of  $M_2$  such that neither  $e$  nor  $f$  is incident with the center of the star  $K_{1,3}$ , then  $G$  is not  $M_2$ -equipackable.*

**Proof.** Without loss of generality, we denote the edges of the star  $K_{1,3}$  by  $vv_1, vv_2, vv_3$ . Neither  $e$  nor  $f$  is incident with  $v$ . For the edge  $e$ , there are three cases:

*Case 1:* The edge  $e$  is not incident with any vertex of  $\{v_1, v_2, v_3\}$ . Since there are at most two vertices of  $\{v_1, v_2, v_3\}$  which are incident with  $f$ , say  $v_1$  and  $v_2$ , we can replace  $\{e, f\}$  with  $\{vv_3, f\}$  and  $\{e, vv_1\}$  to get a maximum  $M_2$ -packing whose size is larger than that of the given maximal  $M_2$ -packing. (See Fig. 1 (a).) So  $G$  is not  $M_2$ -equipackable.

*Case 2:* The edge  $e$  is incident with just one vertex of  $\{v_1, v_2, v_3\}$ , say  $v_1$ . Then  $f$  is not incident with  $v_1$  since  $e$  and  $f$  are independent. We can replace  $\{e, f\}$  with  $\{vv_1, f\}$  and  $\{e, vv_3\}$  to get another larger maximal  $M_2$ -packing which is maximum. (See Fig. 1 (b).) So  $G$  is not  $M_2$ -equipackable.

*Case 3:* The edge  $e$  is incident with two vertices of  $\{v_1, v_2, v_3\}$ , say  $v_1$  and  $v_2$ . Then neither  $v_1$  nor  $v_2$  is incident with  $f$  since  $e$  and  $f$  are independent. We can replace  $\{e, f\}$  with  $\{vv_1, f\}$  and  $\{e, vv_3\}$  to get another larger maximal  $M_2$ -packing. (See Fig. 1 (c).) So  $G$  is not  $M_2$ -equipackable.

In all cases,  $G$  is not  $M_2$ -packable.  $\square$

**Lemma 3.4.** *Let  $G$  be a graph with size  $2m + 1$  and maximum degree  $d$ ,  $2 < d < m + 2$ . If  $G$  is  $M_2$ -equipackable, then for any subgraph  $H$  of  $G$  which is isomorphic to  $K_{1,3}$ ,  $G - H$  is not  $M_2$ -packable.*



**Proof.** Assume that the lemma is not true; that is,  $G - H$  is  $M_2$ -packable. So  $G - H$  can be the union of  $(m - 1)$  copies of  $M_2$ . There exists a copy (say  $\{e, f\}$ ) of  $M_2$  in  $G - H$  such that neither  $e$  nor  $f$  is incident with the center of the star  $H$ . (Otherwise, each copy of  $M_2$  in  $G - H$  has an edge incident with the center (say  $v$ ) of  $H$ , and then  $d = \Delta(G) \geq d_G(v) \geq m - 1 + 3 = m + 2$ , which contradicts that  $d < m + 2$ .) By Lemma 3.3,  $G$  is not  $M_2$ -equipackable. This is a contradiction.  $\square$

**Lemma 3.5.** *Let  $G$  be a graph with size  $2m + 1$ . If there is a maximal  $M_2$ -packing of  $G$  which omits all the edges of a subgraph  $K_3$ , then  $G$  is not  $M_2$ -equipackable.*

**Proof.** Without loss of generality, we denote by  $a, b, c$  the edges of the subgraph  $K_3$ . Select a copy of  $M_2$ , say  $\{e, f\}$ . In the given maximal  $M_2$ -packing of  $G$ , for  $\{e, f\}$ , there are three cases on the adjacency relation with  $\{a, b, c\}$ :

*Case 1:* There is no edge in  $\{e, f\}$  which is adjacent to any edge of  $\{a, b, c\}$ . (See Fig. 2 (a).) We can replace  $\{e, f\}$  with  $\{e, a\}$  and  $\{b, f\}$  to get another  $M_2$ -packing which is maximum. So  $G$  is not  $M_2$ -equipackable.

*Case 2:* Only one edge of  $\{e, f\}$ , say  $e$ , is adjacent to two edges, say  $a, c$ , of  $\{a, b, c\}$ . (See Fig. 2 (b).) Then we can replace  $\{e, f\}$  with  $\{e, b\}$  and  $\{a, f\}$  to get another  $M_2$ -packing which is maximum. So the given  $M_2$ -packing is not maximum and  $G$  is not  $M_2$ -equipackable.

Fig. 1. All cases on the incident relation of  $e$  and  $\{v_1, v_2, v_3\}$ .Fig. 2. All cases on the adjacency relation of  $\{e, f\}$  and  $\{a, b, c\}$ .

*Case 3:* Both of the edges in  $\{e, f\}$  are adjacent to the edges of  $\{a, b, c\}$ . Without loss of generality, suppose that  $e$  is adjacent to  $a, c$  and  $f$  is adjacent to  $b, c$ . (See Fig. 2 (c).) We can replace  $\{e, f\}$  with  $\{e, b\}$  and  $\{a, f\}$  to get another  $M_2$ -packing which is maximum. Thus  $G$  is not  $M_2$ -equipackable. Thus, in all cases,  $G$  is not  $M_2$ -packable.  $\square$

The following corollary easily follows from Lemma 3.5:

**Corollary 3.6.** *Let  $G$  be a graph with size  $2m + 1$ . If  $G$  is  $M_2$ -equipackable, then for any subgraph which is isomorphic to  $K_3$ ,  $G - H$  is not  $M_2$ -packable.*

**Theorem 3.7.** *Let  $G$  be a graph with size  $2m + 1 > 1$  and  $\Delta(G) = d < m + 2$ , then  $G$  is equipackable if and only if  $G$  satisfies one of the following:*

- (1) Neither  $K_3$  nor  $K_{1,3}$  is contained in  $G$  as a subgraph.
- (2) At least one copy of  $K_3$  or  $K_{1,3}$  is contained in  $G$  as a subgraph and, for any subgraph  $H$  of  $G$  which is isomorphic to  $K_3$  or  $K_{1,3}$ ,  $\Delta(G - H) > m - 1$  or  $G - H \cong K_3 \cup K_2$ .

**Proof.** We can easily verify that the graphs described in the theorem are all  $M_2$ -equipackable.

Conversely, let  $G$  be  $M_2$ -equipackable. We consider two cases:

*Case 1:* Neither  $K_3$  nor  $K_{1,3}$  is contained in  $G$  as a subgraph. Clearly,  $G$  is  $M_2$ -equipackable.

*Case 2:* At least one copy of  $K_3$  or  $K_{1,3}$  is contained in  $G$  as a subgraph. By Lemma 3.4 and Corollary 3.6, for any subgraph  $H$  which is isomorphic to  $K_3$  or  $K_{1,3}$ ,  $G - H$  is not  $M_2$ -packable so, by Theorem 1.1,  $\Delta(G - H) > m - 1$  or  $G - H \cong K_3 \cup K_2$ .  $\square$

Especially, we have:

**Corollary 3.8.** *Odd cycles and odd paths are all  $M_2$ -equipackable.*

As far as we know, only  $H$ -equipackable graphs for  $P_3$  and  $M_2$  have been researched. One might consider  $H$ -equipackable graphs for other graphs such as  $P_k$  ( $k = 4, 5, 6$ ),  $M_t$  ( $t \geq 3$ ),  $K_n$  and  $K_{1,r}$ .

## Acknowledgements

We are grateful to Professor P.D. Vestergaard for guiding us to this topic. And we thank Professor Liang Sun for his help. We also wish to thank the referees for prompt, thorough readings of this paper and for many valuable suggestions.

## References

- [1] L.W. Beineke, P. Hamberger, W.D. Goddard, Random packings of graphs, *Discrete Math.* 125 (1994) 45–54.
- [2] B.L. Hartnell, P.D. Vestergaard, Equipackable graphs, *Bulletin of ICA*, R-2005-11.
- [3] J.P. McSorley, T.D. Porter,  $K_p$ -removable sequences of graphs, *J. Combin. Math. Comput.* 45 (2003) 43–62.
- [4] S. Ruiz, Randomly decomposable graphs, *Discrete Math.* 57 (1985) 123–128.
- [6] P.D. Vestergaard, A short update on equipackable graphs, *Discrete Math.*, to appear, *Proceedings of the International Conference Combinatorics'04* in Capomulini, Catania, Italy.